Pis'ma v ZhETF

arXiv:1301.4754v2 [cond-mat.dis-nn] 14 Mar 2013

Ferrimagnetism of dilute Ising antiferromagnets

P. N. Timonin [1]

Southern Federal University, 344090, Rostov-on-Don, Russia

Submitted

It is shown that nearest-neighbor antiferromagnetic interactions of identical Ising spins on imbalanced bipartite lattice and imbalanced bipartite hierarchical fractal result in ferrimagnetic transition instead of antiferromagnetic one. As evidenced by the existing low-field experiments, dilute Ising antiferromagnets may also experience such ferrimagnetic transitions due to the imbalanced nature of the magnetic percolation cluster when it coexists with the percolation cluster of vacancies.

The system of the identical Ising spins on the sites of some crystalline lattices with the nearest-neighbor antiferromagnetic (AF) exchange may have magnetized ground states. In such states there would be antiparallel neighboring spins, as interaction dictates, but the whole numbers of up-spins and down-spins would differ. One such $2d$ lattice is shown in Fig. 1(a). Here two sublattices with parallel up and down spins in the ground state are shown by filled and empty circles correspondingly. We see that in the unit cell there are one filled circle and two empty ones so we get ±1/3 magnetizations in two globally-reversed ground states for nearest-neighbor AF on this lattice. Thus this AF model has a couple of ferrimagnetic ground states with both staggered $L = (\langle S_A \rangle - \langle S_B \rangle)/2$ and homogeneous $M = (2\langle S_A \rangle + \langle S_B \rangle)/3$ magnetizations.

One can easily show that this ordering persists up to finite T_c. Summing the Gibbs function over spins on sublattice A (empty circles) we get the Gibbs distribution for the spins on the sublattice B having effective ferromagnetic Hamiltonian. Indeed, for each link with S_A spin we have (J is AF exchange)

$$\sum_{S_A=\pm 1} \exp\left[-S_A\left(S_B + S'_B J/T\right)\right] = 2\cosh\left[\left(S_B + S'_B J/T\right)\right]$$
$$= 2\exp\left[K_B\left(S_B S'_B + 1\right)\right], \qquad 2K_B \equiv \ln\cosh\left(2J/T\right)$$

Hence, the ordering of S_B spins is described by the ferromagnetic Ising model on the square lattice, so $\langle S_B \rangle \neq 0$ for $K_B > \frac{1}{2}\ln\left(\sqrt{2}+1\right)$ [1] or

$$T < T_c = 2J/\ln\left(\sqrt{2}+1+\sqrt{2\left(\sqrt{2}+1\right)}\right).$$

As $\langle S_B \rangle$ is a linear combination of L and M, the ordered phase is a ferrimagnetic one. This implies that homogeneous magnetic field H has a part conjugated with the order parameter so magnetic susceptibility diverges at T_c and the transition becomes smeared at finite H.



Fig.1. Examples of imbalanced bipartite graphs with different numbers of sites in sublattices A (open circles) and B (filled circles), $N_A > N_B$. (a) - fragment of regular $2d$ lattice, dotted line shows the unit cell; (b, c, d) - clusters of dilute square lattice. In the ground state short-range Ising AF on them would have parallel spins on A and B sublattices and, hence, a nonzero magnetization.

Thus we have a simple example showing that nearest-neighbor AF interaction of identical Ising spins may result in the macroscopic ferrimagnetic order. This is in apparent distinction with conventional ferrimagnets having several different magnetic moments in a cell. It may look rather exotic in the realm of real crystals yet such situation can be frequent in disordered Ising AF, first of all, in dilute Ising AF on bipartite lattices. These lattices can be divided in two subsets of sites, A and B, such that all bonds are of the A-B type, i. e. there are no bonds inside A and B subsets [2]. Apparently, the Ising AF on such lattice is non-frustrated having all spins up on sublattice A and down on sublattice B or vice versa in its two degenerate ground states. Their magnetizations are

$$m = \pm\frac{N_A - N_B}{N_A + N_B} = \pm\frac{1-\eta}{1+\eta},\ \eta \equiv \frac{N_B}{N_A} < 1$$

Here N_A and N_B are the numbers of sites in A and B sublattices and we choose $\eta < 1$ for definiteness.

Seemingly, all known non-frustrated AF crystals with just one sort of magnetic ions have bipartite lattices that are the balanced ones, that is with $\eta = 1$ and purely AF ground states, while Fig. 1(a) shows the imbalanced bipartite lattice with $\eta < 1$ ($\eta = 0.5$). Yet the dilution of balanced bipartite lattices results in appearance of a number of isolated clusters, mostly with $\eta < 1$, those with $\eta = 1$ being the rare exceptions. Figs. 1(b, c, d) show the imbalanced clusters on the square lattice. So at $T = 0$ and arbitrarily small magnetic field dilute bipartite AF must show nonzero magnetization due to the presence of such imbalanced finite clusters. This circumstance was first noticed by Neel [3]. Still it stays unnoticed that for some concentrations of magnetic ions p the giant percolation cluster may also have the average imbalance ratio $\eta_p < 1$.

---

[1] e-mail: pntim@live.ru

Indeed, in finite sample the role of percolation cluster belong to that with the largest number of sites and very probably it is imbalanced, as most of them. However, in the thermodynamic limit ($N \to \infty$) η_p will tend to unity if there are only finite clusters of vacancies. Apparently, such finite clusters cannot make infinite lattice imbalanced as for every cluster deleting unequal number of sites from A and B sublattices there exists (with the same probability) the shifted cluster of the same form which restores the balance. Thus at $1 - p_c < p < 1$ the imbalanced percolation cluster can only exist as a finite-size effect. Meanwhile, at $p_c < p < 1 - p_c$ there is infinite percolation cluster of vacancies to which this argument does not apply. Hence, here $\eta_p < 1$ may also hold in the $N \to \infty$ limit in some crystal lattices. Then the ground state magnetization of dilute AF in this interval will be

$$m_p\left(H = +0\right) = \int_0^1 \frac{1-\eta}{1+\eta} W_p\left(\eta\right) d\eta + \frac{1-\eta_p}{1+\eta_p}$$

Here $W_p\left(\eta\right)$ is the imbalance distribution function of finite clusters. Now it seems that neither $W_p\left(\eta\right)$ nor η_p are known for the crystal lattices. So to find them is quite relevant task for the physics of dilute short-range AF.

The magnetization of finite clusters vanishes at finite temperatures, but that of the percolation cluster could persist up to a finite T_c making the transition a ferrimagnetic one as in the regular imbalanced lattice described above. Then macroscopic features of dilute AF would be those of ordinary ferromagnet in spite of the presence of antiparallel neighboring spins in the ordered phase. In such a case, quite probable for dilute AF in a field (DAFF) at $p_c < p < 1 - p_c$, the mapping of this model onto random-field Ising magnet (RFIM) [4] would be no longer valid. Indeed, the derivation of this correspondence in Ref. [4] is made under the assumption that DAFF has a transition into AF phase which does not hold if, instead, there is a ferrimagnetic transition (smeared at finite H). The evidences in favor of DAFF ferrimagnetism can be found in experiments on several dilute Ising AF with $p < 1 - p_c$ showing the remanent magnetization with the usual order-parameter behavior [5]-[7] and prominent peak in temperature dependence of magnetic susceptibility which appears in low fields as a result of dilution and becomes gradually smeared in higher fields [8],[9].

To get some notion of the thermodynamics of possible DAFF ferrimagnetism we consider here the nearest-neighbor AF on the simplest hierarchical lattice, imitating the percolation cluster with fractal dimension $d = 2$ and $\eta = 1/3$. It may describe qualitatively dilute AF, large planar aggregates of AF particles or disordered AF thin films. The model also exhibits a number of field-induced ground state transitions marked by the magnetization jumps which are characteristic for dilute AF [10], [11].

LOW-FIELD THERMODYNAMICS OF HIERARCHICAL ANTIFERROMAGNET

We consider the short-range Ising AF on the simplest "diamond" hierarchical lattice [12]. Its building process is shown in Fig.2. On the n-th level of hierarchy the lattice has N_n sites, $N_n = \frac{2}{3}\left(4^n + 2\right)$, see Ref. [13]. The coordination numbers of the sites are the powers of 2: $z = 2, 4, 8, \ldots$. At all levels of the hierarchy the lattice is bipartite and for $n > 0$ the sites with coordination number $z = 2$ constitutes the sublattice A (open circles in Fig.2) while the others belong to the sublattice B (filled circles), $N_{A,n} \geqslant N_{B,n}$. At the n-th level $N_{A,n} = 2 \cdot 4^{n-1}$ for $n > 0$ [13], so $\eta_n = (N_n - N_{A,n})/N_{A,n} = \left(1 + 2 \cdot 4^{1-n}\right)/3$. We are interested in the thermodynamic limit of infinite levels of hierarchy. In this limit $\eta = 1/3$ and fractal dimension $d = 2$ [13].



Fig.2. Construction of hierarchical lattice. It is bipartite at all levels. Different circles designate its partitioning, open circles sublattice A, filled circles sublattice B, $N_A \geq N_B$.

For the Ising spins $S_i = \pm 1$ placed on the sites of this lattice we consider the AF Hamiltonian

$$\mathcal{H} = J \sum_{<i\in A, j\in B>} S_i S_j - H_A \sum_{i\in A} S_i - H_B \sum_{j\in B} S_j \qquad (1)$$

where $< i \in A, j \in B >$ means the summation over nearest neighbors and different fields for the sublattices are introduced. This allows to find the average magnetizations of each sublattice and the order parameter for the transition. Homogeneous field corresponds to $H_A = H_B = H$.

The usual way to get the partition function of the model is through the recursion relations for partial partition functions at different levels of hierarchy $Z_n(S, S')$ having fixed values of the outmost left and right spins S and S' [12]. These relations read

$$Z_{n+1}(S,S') = \left[\sum_{S_1=\pm 1} Z_n(S,S_1) e^{h_n S_1} Z_n(S_1,S') \right]^2, \qquad (2)$$

$h_n = H_n/T$, $H_0 = H_A$ and $H_n = H_B$, $n \geqslant 1$. The initial condition for them is

$$Z_0(S,S') = e^{-KSS'}, \qquad K = J/T. \qquad (3)$$

Using the representation

$$Z_n(S,S') = \exp \frac{1}{2} \left[C_n + u_n S S' + (v_n - h_n)(S + S') \right]$$

we get from Eqs.(2,3)

$$\begin{aligned} u_0 = -2K, \qquad v_0 = h_A, \qquad C_0 = 0 \\ u_{n+1} = 2 \ln \cosh u_n + \ln \left(1 - \tanh^2 u_n \tanh^2 v_n \right), \\ v_{n+1} = 2v_n + 2\tanh^{-1}(\tanh u_n \tanh v_n) - 2h_n + h_{n+1}, \\ C_{n+1} = 4C_n + u_{n+1} + 4\ln(2\cosh v_n). \qquad (4) \end{aligned}$$

The last of Eqs.(4) gives for $n > 0$

$$C_n = u_n - 4^n u_0 + \sum_{l=0}^{n-1} 4^{n-l} \left[u_l + \ln(2\cosh v_l) \right]$$

so the n-th level free energy per spin is

$$F_n = -\frac{T}{N_n} \ln \sum_{S,S'} Z_n(S,S') e^{h_B(S+S')} = -\frac{3}{4} T \sum_{l=0}^{n-1} 4^{-l} \left[u_l + \ln(2\cosh v_l) \right] - \frac{3}{2} J + O(1/N_n) \qquad (5)$$

At $H_n = 0$ the model has phase transition at $K = K_c \approx 0.609$ being the solution to the equation $K_c = \ln\cosh 2K_c$. $u_c = 2K_c$ is the stationary point of the zero-field equations, $u_{n+1} = 2\ln\cosh u_n$, $v_n = 0$. In the paramagnetic phase at $K < K_c$ $u_n \to 0$ for $n \to \infty$, while in the ordered phase at $K > K_c$ $u_n \to \infty$. According to above considerations the ground states of the model have magnetizations $\pm(1/2)$, so we may expect that the ordered phase is ferrimagnetic. To show this we consider Eqs.(4) at

$$0 < (T_c - T)/T_c \equiv \tau \ll 1, \qquad |h_n| \ll \tau. \qquad (6)$$

In this case u_n and v_n can be found approximately in the three regions of n:

1) $1 \leqslant n \leqslant \lambda, u_n - u_c \leqslant u_c, |v_n| \ll 1$

$$u_n \approx u_c + \kappa^{n-1}(u_1 - u_c), \qquad (7)$$

$$v_n \approx \frac{h_B + (2+\kappa)^{n-1} \kappa \tilde{h}}{1+\kappa}, \qquad \tilde{h} = h_B - (\kappa+1) h_A \qquad (8)$$

$$u_1 = 2\ln\cosh 2K, \qquad \kappa = 2\tanh 2K_c \approx 1.68,$$

The value of λ is defined by

$$u_\lambda = 2u_c, \qquad \kappa^{-\lambda} = \frac{u_1 - u_c}{\kappa u_c} \approx \tau \qquad (9)$$

while $|v_n| \ll 1$ requires

$$|\tilde{h}|(2+\kappa)^\lambda = |\tilde{h}| \tau^{-\ln(2+\kappa)/\ln\kappa} \ll 1. \qquad (10)$$

2) $\lambda < n \leqslant \mu, 1 \ll u_n \ll |v_n|$

$$u_n \approx 2^{n-\lambda+1} u_c - \sum_{k=\lambda}^{n-1} 2^{n-k} \ln 2\cosh v_k, \qquad (11)$$

$$v_n \approx \frac{h_B}{3} + 4^{n-\lambda} \left(v_\lambda - \frac{h_B}{3} \right), \qquad (12)$$

$$v_\lambda \approx \frac{(2+\kappa)^{\lambda-1} \kappa \tilde{h}}{(1+\kappa)} = \frac{\kappa \tilde{h}}{(1+\kappa)(2+\kappa)\tau^{\ln(2+\kappa)/\ln\kappa}} \qquad (13)$$

The value of μ is defined by the equation $u_\mu = |v_\mu|$. As $u_\mu \approx 2^{\mu-\lambda+1} u_c - \sum_{k=\lambda}^{\mu-1} 2^{\mu-k} \ln 2\cosh 4^{k-\lambda} v_\lambda \approx 2^{\mu-\lambda} 2(u_c - \ln 2) + 4^{\mu-\lambda} v_\lambda$, $v_\mu \approx 4^{\mu-\lambda} v_\lambda$, we get

$$2^\mu = \frac{(u_c - \ln 2) 2^\lambda}{|v_\lambda|}, \quad u_\mu = |v_\mu| = \frac{(u_c - \ln 2)^2}{|v_\lambda|}. \qquad (14)$$

$$3) \mu > n, |v_n| \gg 1, u_n \approx 0, v_n \approx 2^{n-\mu} v_\mu. \qquad (15)$$

Note that in the sums we consider the large numbers λ and μ as integers neglecting its fractional parts.

Using the above approximations for u_n and v_n we can find from Eq.(5) free energy in the thermodynamic limit near the transition point in a small field (cf.

Eqs.(6), (10)). Thus, dividing the sum in (5) in three parts and $n=0$ term,

$$-\frac{4}{3}\frac{F}{T}=\ln 2\cosh h_A+\Sigma_\lambda+\Sigma_{\lambda\mu}+\Sigma_\mu,$$

we get

$$\begin{aligned}\Sigma_\lambda=&\sum_{n=1}^{\lambda}\left[4^{-n}\left(u_c+\ln 2\right)+u_c\tau\left(\frac{\kappa}{4}\right)^n\right]+\\&+\sum_{n=1}^{\lambda}\frac{\tilde{h}^2}{2(1+\kappa)^2(2+\kappa)^2}\left(1+\frac{\kappa}{2}\right)^{2n}\\\approx&\frac{1}{3}\left(\ln 2+u_c\right)+\frac{\kappa}{4-\kappa}u_c\tau-\frac{4^{-\lambda}}{3}\left(\frac{2+\kappa}{4-\kappa}2u_c+\ln 2\right)\\&+\frac{\kappa\tilde{h}^2}{2\left(4+\kappa\right)\left(1+\kappa\right)^2}\left(1+\frac{\kappa}{2}\right)^{2\lambda}\end{aligned}$$

$$\begin{aligned}\Sigma_{\lambda\mu}=&\sum_{n=\lambda+1}^{\mu}\left(2^{-n-\lambda+1}u_c+4^{-n}\ln 2\cosh v_n\right)-\\&-\sum_{n=\lambda+1}^{\mu}4^{-n}\sum_{k=\lambda}^{n-1}2^{n-k}\ln 2\cosh v_k\\=&\sum_{n=\lambda+1}^{\mu}\left(2^{-n-\lambda+1}u_c+4^{-n}\ln 2\cosh v_n\right)-\\&-\sum_{k=\lambda}^{\mu}\left(4^{-k}-2^{-k-\mu}\right)\ln 2\cosh v_k\\=&4^{-\lambda}2u_c-4^{-\mu}u_\mu+4^{-\mu}\ln 2\cosh v_\mu-4^{-\lambda}\ln 2\cosh v_\lambda\\\approx&4^{-\lambda}\left(2u_c-\ln 2\right)\end{aligned}$$

$$\Sigma_\mu\approx\sum_{n=\mu+1}^{\infty}4^{-n}\left|v_n\right|\approx 4^{-\mu}\left|v_\mu\right|=4^{-\lambda}\left|v_\lambda\right|.$$

Here we used $|v_\lambda|\ll 1,\ |v_\mu|\gg 1$, Eqs.(7, 8, 11, 12, 15) and the relation following from Eq.(11),

$$\sum_{k=\lambda}^{\mu-1}2^{-k-\mu}\ln 2\cosh v_k=2^{-\lambda-\mu+1}u_c-4^{-\mu}u_\mu.$$

Finally we have from Eqs.(5, 9, 13, 14)

$$F/T_c\approx -K_c/2-\ln 2+\tau s_c-a\tau^{2-\alpha}-b\tau^{\beta}\left|\tilde{h}\right|-c\tau^{-\gamma}\tilde{h}^2,\tag{16}$$

$s_c=\ln 2-2K_c\frac{\kappa-1}{4-\kappa}\approx 0.34$, $a=2K_c\frac{5-2\kappa}{4-\kappa}-\ln 2\approx 0.17$, $b=\frac{3\kappa}{4(2+\kappa)(1+\kappa)}\approx 0.13$, $c=\frac{3\kappa}{8(4+\kappa)(1+\kappa)^2}\approx 0.015$.

$$\begin{aligned}\alpha=2-\frac{\ln 4}{\ln\kappa}\approx -0.67,\beta=\frac{\ln 4-\ln\left(2+\kappa\right)}{\ln\kappa}\approx 0.16,\\\gamma=\frac{2\ln\left(2+\kappa\right)-\ln 4}{\ln\kappa}\approx 2.35\end{aligned}\tag{17}$$

In homogeneous field $\tilde{h}=-\kappa h$ (cf. Eq.(8)) so F in Eq.(16) has the standard scaling form of a ferromagnet with spontaneous magnetization $m\sim\tau^{\beta}$ and divergent susceptibility $\chi\sim\tau^{-\gamma}$. This expression is valid at $0<\tau\ll 1$, $|h|\ll\tau^{\beta+\gamma}$, cf. Eq.(10). Scaling indices (17) obey the usual relation $\alpha+2\beta+\gamma=2$. Negative α means that specific heat is finite at the transition point and has a cusp at T_c . Note also that s_c is the entropy at the transition point. So this AF system looks like genuine ferromagnet, even featuring the absence (smearing) of transition in a finite field. The last is evident as the nontrivial stationary point of finite-field recursion relations (4) cannot be reached from any initial conditions.

Yet the dependence of F on $\tilde{h}$ from Eq.(8) shows that true order parameter for the transition is a linear combination of $M_A=\sum\limits_{i\in A}S_i$ and $M_B=\sum\limits_{i\in B}S_i$ conjugate with $\tilde{H}=H_B-\left(\kappa+1\right)H_A$. To distinguish the order parameter in the Hamiltonian (1) we perform a coordinate rotation in $2d$ space of vectors $\mathbf{H}=\left(H_A,H_B\right)$ and $\mathbf{M}=\left(M_A,M_B\right)$ to bring the term $-\mathbf{MH}$ in (1) to the form $-\mathbf{MH}=-\tilde{M}\tilde{H}-M'H'$ where

$$\tilde{M}=\frac{M_B-\left(\kappa+1\right)M_A}{1+\left(\kappa+1\right)^2},M'=\frac{\left(\kappa+1\right)M_B+M_A}{1+\left(\kappa+1\right)^2},$$
$$H'=\left(\kappa+1\right)H_B+H_A$$

Thus $\tilde{M}$ is the order parameter while M' and H' are non-critical variables. Hence, $\langle M'\rangle=0$ at $\boldsymbol{H}\to 0$ so the spontaneous magnetic moments of the sublattices obey the relation $\langle M_A\rangle=-\left(\kappa+1\right)\langle M_B\rangle$. Then for the spontaneous magnetizations $m_\nu=\langle M_\nu\rangle/N_\nu$ ($\nu=\ A,\ B$) we have

$$m_A=-\eta\left(\kappa+1\right)m_B\approx -0.9m_B$$

This differs from the ground state relation $m_A=-m_B$. We may suggest that this is a consequence of critical fluctuations diminishing m_A more strongly than m_B as all sites of sublattice A have the lowest coordination number $z_A=2$. To some extent this effect would be present in all dilute AF on imbalanced bipartite graphs since the sublattice A with larger amount of spins would necessary have lower average coordination number $\bar{z}_A=C/N_A<\bar{z}_B=C/N_B$. Here C is the number of bonds and we used the fact that all bonds are of A-B type. Thus $\bar{z}_A=\eta\bar{z}_B$ so the lower η the stronger can be the fluctuation-induced disbalance between m_A and m_B near T_c. At $\eta=1$ this effect vanishes so its observation in neutron-diffraction experiments can certify the onset of imbalance in the magnetic percolation cluster.

GROUND STATE TRANSITIONS

Here we assume $H_A = H_B = H$. At $T = 0$ we define $\tilde{u}_n = \lim_{T\to 0} T u_n$, $\tilde{v}_n = \lim_{T\to 0} T v_n$, $E = \lim_{T\to 0} F$ to obtain from Eqs.(4, 5)

$$\tilde{u}_0 = -2J \qquad \tilde{v}_0 = H$$

$$\tilde{u}_{n+1} = 2\left(|\tilde{u}_n| - |\tilde{v}_n|\right)\vartheta\left(|\tilde{u}_n| - |\tilde{v}_n|\right), \tag{18}$$

$$\tilde{v}_{n+1} = 2\tilde{v}_n + 2\min\left(|\tilde{u_n}|, |\tilde{v_n}|\right)\operatorname{sgn}\left(\tilde{u}_n\tilde{v}_n\right) - H, \tag{19}$$

$$-(4/3)E = H + \sum_{n=1}^{\infty} 4^{-n}\left(\tilde{u}_n + |\tilde{v}_n|\right)$$

ϑ in Eq.(18) is Heaviside's step function. Solving Eqs.(18, 19), we get

$$-(4/3)E = H + \sum_{n=1}^{\infty} 4^{-n}\left|H - 2^{n+1}J\right|,$$

$$m = -\frac{\partial E}{\partial H} = \frac{3}{4}\left[1 + \sum_{n=1}^{\infty} 4^{-n}\operatorname{sgn}\left(H - 2^{n+1}J\right)\right].$$

So at $H \neq 2^k J$

$$m = 2^{-1}\vartheta\left(2J - H\right) + \left(1 - 2\cdot 4^{-r}\right)\vartheta\left(H - 2J\right),$$

where $r = [\log_2(H/J)]$ is an integer part of $\log_2(H/J)$. At $H_r = 2^r J$, $r \geqslant 2$, we have $m_r = 1 - 6\cdot 4^{-r}$. Field dependence of the ground state magnetization is shown in Fig.3. Due to the imbalance ratio $\eta = 1/3$ the system has spontaneous magnetization $m = 1/2$ at $H \to +0$. The jumps at $H_r = 2^r J$ result from the flipping along the field of single spins in sublattice B having the coordination number 2^r. Note that in dilute crystalline lattices there are many more jumps appearing at rational values of H/J where flipping of the whole clusters takes place [10]. Such jumps were observed in low-T experiments in $Fe_pZn_{1-p}F_2$ [11]. The ground states of DAFF on cubic and bcc lattices are studied numerically for all p in Ref. [14]. Unfortunately, the data on the sample's and percolation cluster magnetization are totally absent in this paper so we cannot decide if there is the imbalance in real $3d$ percolation clusters at $p_c < p < 1 - p_c$.



Fig.3. Field dependence of the ground state magnetization.

DISCUSSION AND CONCLUSIONS

The decades of experimental investigations of dilute Ising AF have shown that DAFF - RFIM correspondence works reasonably well at low dilution [15]. Meanwhile the field-induced rounding of the transition appears at lower p which is impossible in the case of the AF ordered phase. One explanation assumes that this is nonequilibrium effect due to the pinning of AF domain walls by the vacancies which results in very slow relaxation to the equilibrium AF structure [16]. The numerical simulations demonstrate that this effect is strongly enhanced with the appearance of vacancies' percolation [17].

Yet the experiments show that not just a slowing down but true nonergodic irreversibility with multiple metastable states sets in at intermediate p [18], so one may suggest that AF transition transforms into a spin-glass one [15],[19]. This is quite probable scenario at p close to p_c where weak next-nearest interactions may cause a frustration in a sparse percolation cluster. However, the field-induced rounding and the nonergodic irreversibility sets in right below $p = 1 - p_c$ in real dilute AF [20]. The observation of Ref. [20] that AF transition breaks up with the appearance of vacancies' percolation, along with the ferromagnetic anomalies found in Refs. [5]-[9] at $p_c < p < 1 - p_c$, strongly supports the onset of imbalance in magnetic percolation cluster making the transition ferrimagnetic in this concentration range. This explains the field-induced rounding, while the weak dipole-dipole interactions would give rise to a number of metastable magnetized domain states thus explaining the appearing nonergodicity. The expected ferrimagnetic phase at intermediate dilution would differ macroscopically from the spin-glass one by a small scaling index of remanent magnetization $\beta < 0.5$ and by the divergent magnetic susceptibility. The present model also predicts rather high index γ, large negative α and disbalance in the sublattice magnetizations near T_c. But to observe these features of ferrimagnetic transition the measurements in ultra-low fields (same as in Refs. [5]-[9]) are needed to avoid the smearing of the transition.

Here we can only give qualitative arguments in favor of ferrimagnetism of real dilute AF, explaining the results of Refs. [5]-[9],[18],[20]. Now we do not know how to rigorously prove that percolation cluster on dilute $3d$ lattices has $\eta_p < 1$ at $p_c < p < 1 - p_c$. Yet this hypothesis can be verified by numerical simulations. This and further low-field experiments may help to elucidate the nature of transition in nearest-neighbor dilute AF.

Author gratefully acknowledges useful discussions with V.P. Sakhnenko.

1. R. J. Baxter, *Exactly solved models in statistical mechanics*, Academic Press, 1982.
2. G. Chatrand, *Introductory Graph Theory*, Dover, 1984.
3. L. Neel, C. R. Acad. Sci. Paris **252**, 4075 (1961).
4. J. L. Cardy, Phys. Rev. **B29**, R2460 (1984).
5. C. Djurberg, J. Mattson and P. Nordblad, J. Appl. Phys. **75**, 5541 (1994); J. Mattson, C. Djurberg and P. Nordblad Phys. Rev. **B61**, 11274 (2000).
6. J. Kushauer, W. Kleemann, J. Mattsson et al., Phys. Rev. **B49**, 6346 (1994).
7. M. Lederman, J. Hammann and R. Orbach, Physica **B165-166**, 179 (1990).
8. H. Ikeda, J. Phys. C **16**, L21 (1983).
9. H. Ikeda and K. Kikuta, J. Phys. C **16**, L445 (1983).
10. S. Bastea and P. M. Duxbury, Phys. Rev. **E58**, 4261 (1998)
11. A. R. King, V. Jaccarino, T. Sakakibara et al., Phys. Rev. Lett. **47**, 117 (1981).
12. A. N. Berker and S. Ostlund, J. Phys. C **12**, 4961 (1979); R. B. Griffiths and M. Kaufman, Phys. Rev. **B26**, R5022 (1982).
13. P. N. Timonin, Zh. Eksp. Teor. Fiz. **126**, 1198 (2004).
14. A. Glaser, A. C. Jones and P. M. Duxbury, Phys. Rev. **B71**, 174423 (2005).
15. D. P. Belanger in *Spin Glasses and Random Fields*, ed. A. P. Young, World Scientific (1997).
16. M. Staats, U. Nowak and K. D. Usadel, Phase Transitions **65**, 159 (1998).
17. W. C. Barber and D. P. Belanger, J. Appl. Phys. **87**, 7049 (2000).
18. M. Lederman, J. V.Seilinger, R. Bruinsma et al., Phys. Rev. **B48**, 3810 (1993).
19. F. C. Montenegro, S. M. Rezende and M. D. Coutinho-Filho, J. Appl. Phys., **63**, 3755 (1988).
20. W. C. Barber, F. Ye, D. P. Belanger et al., Phys. Rev. **B69**, 024409 (2004).